

Q^2 evolution of parton distributions at low x . Soft initial conditions

A.V. Kotikov

*Particle Physics Laboratory
Joint Institute for Nuclear Research
141980 Dubna, Russia*

G. Parente

*Departamento de Física de Partículas
Universidade de Santiago de Compostela
15706 Santiago de Compostela, Spain*

Abstract

We investigate the Q^2 evolution of parton distributions at small x values, recently obtained in the case of soft initial conditions. The results are in excellent agreement with deep inelastic scattering experimental data from HERA.

The measurements of the deep-inelastic scattering structure function F_2 in HERA [1] have permitted the access to a very interesting kinematical range for testing the theoretical ideas on the behavior of quarks and gluons carrying a very low fraction of momentum of the proton, the so-called small x region. In this limit one expects that non-perturbative effects may give essential contributions. However, the reasonable agreement between HERA data and the NLO approximation of perturbative QCD that has been observed for $Q^2 > 1\text{GeV}^2$ (see the recent review in [2]) indicates that perturbative QCD could describe the evolution of structure functions up to very low Q^2 values, traditionally explained by soft processes. It is of fundamental importance to find out the kinematical region where the well-established perturbative QCD formalism can be safely applied at small x .

The standard program to study the small x behavior of quarks and gluons is carried out by comparison of data with the numerical solution of the DGLAP equations ¹ by fitting the parameters of the x profile of partons at some initial Q_0^2 and the QCD energy scale Λ (see, for example, [3]). However, if one is interested in analyzing exclusively the small x region, there is the alternative of doing a simpler analysis by using some of the existing analytical solutions of DGLAP in the small x limit (see [2] for review). This was done so in Ref. [4]-[6] where it was pointed out that the HERA small x data can be interpreted in terms of the so called doubled asymptotic scaling phenomenon related to the asymptotic behaviour of the DGLAP evolution discovered in [7] many years ago. Here we will illustrate results obtained recently in [6].

1. Thus, our purpose is to demonstrate the small x asymptotic form of parton distributions in the framework of the DGLAP equation starting at some Q_0^2 with the flat function:

$$f_a(Q_0^2) = A_a \quad (a = q, g), \quad (1)$$

¹ At small x there is another approach based on the BFKL equation, whose application is out of the scope

where f_a are the parton distributions multiplied by x and A_a are unknown parameters that have to be determined from data. Through this work at small x we neglect the non-singlet quark component.

In [6] an effective method to reproduce the x -dependence of parton distributions has been developed. It is based on a separation of the singular and regular parts of the exact solution for the moments of parton distributions. The method allows in simplest way to reproduce the leading order (LO) results [4] and to construct the x -dependence of parton distributions at next-to-leading order (NLO):

$$\begin{aligned}
f_a(x, Q^2) &= f_a^+(x, Q^2) + f_a^-(x, Q^2) \quad \text{and} \\
f_a^-(x, Q^2) &= A_a^-(Q^2, Q_0^2) \exp(-d_-(1)s - D_-(1)p) + O(x) \\
f_g^+(x, Q^2) &= A_g^+(Q^2, Q_0^2) I_0(\sigma) \exp(-\bar{d}_+(1)s - \bar{D}_+(1)p) + O(\rho) \\
f_q^+(x, Q^2) &= A_q^+(Q^2, Q_0^2) \left[(1 - \bar{d}_\pm^q(1)\alpha(Q^2)) \rho I_1(\sigma) + 20\alpha(Q^2) I_0(\sigma) \right] \\
&\quad \cdot \exp(-\bar{d}_+(1)s - \bar{D}_+(1)p) + O(\rho) \\
F_2(x, Q^2) &= e \cdot \left(f_q(x, Q^2) + \frac{2}{3} f \alpha(Q^2) f_g(x, Q^2) \right)
\end{aligned} \tag{2}$$

where $I_\nu(\sigma)$ is modified Bessel function, s and p are given by $s = \ln(\alpha(Q_0^2)/\alpha(Q^2))$, $p = \alpha(Q_0^2) - \alpha(Q^2)$, and

$$\sigma = 2\sqrt{(\hat{d}_+ s + \hat{D}_+ p) \ln x} \quad , \quad \rho = \sqrt{\frac{(\hat{d}_+ s + \hat{D}_+ p)}{\ln x}} = \frac{\sigma}{2\ln(1/x)}$$

are NLO generalizations of the corresponding Ball-Forte variables and

$$\begin{aligned}
A_g^+(Q^2, Q_0^2) &= \left[1 - \frac{80}{81} f \alpha(Q^2) \right] A_g + \frac{4}{9} \left[1 + 3 \left(1 + \frac{1}{81} f \right) \alpha(Q_0^2) - \frac{80}{81} f \alpha(Q^2) \right] A_q \quad , \\
A_g^-(Q^2, Q_0^2) &= A_g - A_g^+(Q^2, Q_0^2) \\
A_q^+ &= \frac{f}{9} \left(A_g + \frac{4}{9} A_q \right) \quad , \quad A_q^- = A_q - 20\alpha(Q_0^2) A_q^+
\end{aligned} \tag{3}$$

Wherever in this work we use the notation $\alpha = \alpha_s/(4\pi)$.

The nonzero components of the singular and regular parts of the terms $d_\pm$ and $D_\pm$ have the form:

$$\begin{aligned}
\hat{d}_+ &= -\frac{12}{\beta_0} \quad , \quad \bar{d}_+(1) = 1 + \frac{4}{3\beta_0} f \quad , \quad d_-(1) = \frac{16}{27\beta_0} f \quad , \\
\hat{d}_{++} &= \frac{412}{27\beta_0} f \quad , \quad \hat{d}_{+-}^q = -20 \quad , \quad \bar{d}_{+-}^q(1) = \frac{80}{81} f \quad , \\
\bar{d}_{++}(1) &= \frac{8}{\beta_0} \left(36\zeta_3 + 33\zeta_2 - \frac{1643}{12} + \frac{2}{9} f \left[\frac{68}{9} - 4\zeta_2 - \frac{13}{243} f \right] \right) \quad , \\
\bar{d}_{+-}^q(1) &= \frac{134}{3} - 12\zeta_2 - \frac{13}{81} f \quad , \quad \bar{d}_{-+}^q(1) = -3 \left(1 + \frac{f}{81} \right) , \\
d_{--}(1) &= \frac{16}{9\beta_0} \left(2\zeta_3 - 3\zeta_2 + \frac{13}{4} + f \left[4\zeta_2 - \frac{23}{18} + \frac{13}{243} f \right] \right) \quad ,
\end{aligned} \tag{4}$$

where β_0 is the first coefficient of QCD β -function, ζ_n are Euler ζ -functions and f is the number of active quarks.

- Our NLO results coincide with the corresponding of Ball and Forte in Ref. [5] if one neglects the “−” component, expands our NLO singular terms $(\rho)^k I_{k+1}(\sigma)$ in the vicinity of the point $\sigma = \sigma_{LO}$ and ignores the NLO regular terms (i.e. put $\exp(-\overline{D}_+(1)p) = \exp(-D_-(1)p) = 1$ and cancel the terms proportional to $\alpha(Q_0^2)$ into the normalization factors $A_g^\pm$ and $A_q^\pm$). We think, however, that this expansion is not so correct because it generates NLO corrections of the order of the LO terms.
- The behaviour of eqs. (2) can mimic a power law shape over a limited region of x, Q^2 :

$$f_a(x, Q^2) \sim x^{-\lambda_a^{eff}(x, Q^2)} \quad \text{and} \quad F_2(x, Q^2) \sim x^{-\lambda_{F_2}^{eff}(x, Q^2)}$$

The quark and gluon effective slopes λ_a^{eff} are reduced by the NLO terms that leads to the decreasing of the gluon distribution at small x . For the quark case it is not the case, because the normalization factor A_q^+ of the “+” component produces an additional contribution undampening as $\sim (\ln x)^{-1}$.

- The gluon effective slope λ_g^{eff} is larger than the quark slope λ_q^{eff} , which is in excellent agreement with a recent MRS and GRV analyses [3].
Indeed, the effective slopes have the asymptotical values (at large Q^2):

$$\begin{aligned} \lambda_g^{eff,as}(x, Q^2) &= \rho \frac{I_1(\sigma)}{I_0(\sigma)} \approx \rho - \frac{1}{4 \ln(1/x)} \\ \lambda_q^{eff,as}(x, Q^2) &= \rho \cdot \frac{I_2(\sigma)(1 - \overline{d}_{+-}^q(1)\alpha(Q^2)) + 20\alpha(Q^2)I_1(\sigma)/\rho}{I_1(\sigma)(1 - \overline{d}_{+-}^q(1)\alpha(Q^2)) + 20\alpha(Q^2)I_0(\sigma)/\rho} \\ &\approx \left(\rho - \frac{3}{4 \ln(1/x)} \right) \left(1 - \frac{10\alpha(Q^2)}{(\hat{d}_{+s} + \hat{D}_{+p})} \right) \\ \lambda_{F_2}^{eff,as}(x, Q^2) &= \lambda_q^{eff,as}(x, Q^2) \frac{1 + 6\alpha(Q^2)/\lambda_q^{eff,as}(z, Q^2)}{1 + 6\alpha(Q^2)/\lambda_g^{eff,as}(z, Q^2)} + O(\alpha^2(Q^2)) \\ &\approx \lambda_q^{eff,as}(z, Q^2) + \frac{3\alpha(Q^2)}{\ln(1/x)}, \end{aligned} \tag{5}$$

where symbol $\approx$ marks approximations obtained by expansions of modified Bessel functions $I_n(\sigma)$. The slope $\lambda_{F_2}^{eff,as}(x, Q^2)$ lies between quark and gluon ones but closely to quark slope $\lambda_q^{eff,as}(x, Q^2)$ (see also Fig. 2).

- Both slopes λ_a^{eff} decrease with decreasing x . A x dependence of the slope should not appear for a PD with a Regge type asymptotic ($x^{-\lambda}$) and precise measurement of the slope λ_a^{eff} may lead to the possibility to verify the type of small x asymptotics of parton distributions.

2. With the help of the results obtained in the previous section we have analyzed F_2 HERA data at small x from the H1 collaboration (first article in [1]). In order to keep the analysis as simple as possible we have fixed $\Lambda_{\overline{\text{MS}}}(n_f = 4) = 250$ MeV which is a reasonable value extracted from the traditional (higher x) experiments. The initial scale of the PD was also fixed into the fits to $Q_0^2 = 1 \text{ GeV}^2$, although later it was released to study the sensitivity of the fit to the variation of this parameter. The analyzed data region was restricted to $x < 0.01$ to remain within the kinematical range where our results are accurate. Finally, the number of active flavors was fixed to $f=4$.

Fig. 1 shows F_2 calculated from the fit with $Q^2 > 1 \text{ GeV}^2$ in comparison with H1 data. Only the data with Q^2 higher than Q_0^2 are shown, so that the NLO result (dotted-dashed line) lies

closer to the data than the LO curve (dashed line). The lack of agreement between data and lines observed at the lowest x and Q^2 bins suggests that the flat behavior should occur at Q^2 lower than 1 GeV^2 . In order to study this point we have done the analysis considering Q_0^2 as a free parameter. Comparing the results of the fits (see [6]) one can notice the better agreement with the experiment of the NLO curve at fitted $Q_0^2 = 0.55 \text{ GeV}^2$ (solid curve) is apparent at the lowest kinematical bins.

Finally with the help of Eqs. (5) we have estimated the F_2 effective slope using the value of the parameters extracted from NLO fits to data. For H1 data we found $0.05 < \lambda_{F_2}^{eff} < 0.30 - 0.37$. The lower (upper) limits corresponds ² to $Q^2 = 1.5 \text{ GeV}^2$ ($Q^2 = 400 \text{ GeV}^2$). The dispersion in some of the limits is due to the x dependence. Fig. 2 shows that the three types of asymptotical slopes have similar values, which are in very good agreement with H1 data (presented also in Fig. 2). The NLO values of $\lambda_{F_2}^{eff,as}$ lie between the quark and the gluon ones but closer to the quark slope $\lambda_q^{eff,as}$. These results are in excellent agreement with those obtained by others (see the review [2] and references therein).

3. Resume. We have shown that the results developed recently in [6] have quite simple form and reproduce many properties of parton distributions at small x , that have been known from global fits.

We found very good agreement between our approach based on QCD at NLO approximation and HERA data, as it has been observed earlier with other approaches (see the review [2]). Thus, the nonperturbative contributions as shadowing effects, higher twist effects and others seems to be quite small or seems to be canceled between them and/or with $\ln(1/x)$ terms containing by higher orders of perturbative theory. To clear up the correct contributions of nonperturbative dynamics and higher orders containing strong $\ln(1/x)$ terms, it is necessary more precise data and further efforts in developing of theoretical approaches.

One of the authors (G.P.) was supported in part by Xunta de Galicia (XUGA-20604A96) and CICYT (AEN96-1673).

References

- [1] H1 Collab.:S. Aid et al., *Nucl.Phys.* **B470** (1996) 3; ZEUS Collab.: M. Derrick et al., *Zeit.Phys.* **C72** (1996) 399; **C69** (1996) 607.
- [2] A. M. Cooper-Sarkar et al., *Int.J.Mod..Phys.* **A13** (1998) 3385.
- [3] A.D. Martin, W.S. Stirling and R.G. Roberts, *Phys.Lett.* **B387** (1996) 419; M. Gluck, E. Reya and A. Vogt, D0-TH 98/07, WUE-ITP-98-019 (hep-ph/9806404).
- [4] R.D. Ball and S. Forte, *Phys.Lett.* **B336** (1994) 77; L. Mankiewicz, A. Saalfeld and T. Weigl, *Phys.Lett.* **B393** (1997) 175.
- [5] R.D. Ball and S. Forte, *Acta Phys.Polon.* **B26** (1995) 2097; *Nucl. Phys. Proc. Suppl.* **54A** (1997) 163.
- [6] A.V. Kotikov and G. Parente, Preprint US-FT/13-98 (hep-ph/9807249).
- [7] A. De Rújula et al., *Phys.Rev.* **D10** (1974) 1649.

² $\lambda_{F_2}^{eff} = \frac{1}{Q^2} \frac{dF_2}{d \ln Q^2}$ and the same for the other slopes.

Fig. 1




Fig. 2

